

Full Length Research Paper

Micropropagation of *Launaea cornuta* - an important indigenous vegetable and medicinal plant

Faith Ambajo¹ and Jonathan Mutie Matheka^{2*}

¹Department of Biotechnology, University of Eldoret, P.O. Box 1125, Eldoret, Kenya.

²International Institute of Tropical Agriculture c/o ICIPE P.O. Box 30772-00100 Nairobi, Kenya.

Received 6 January, 2016; Accepted 19 July, 2016

An efficient micropropagation protocol was developed for the medicinal plant *Launaea cornuta* using green house axillary buds as explants. The best sterility was obtained at 30% (v/v) local bleach (JIK). Maximum shoot induction rate was achieved when axillary buds were cultured on Murashige and Skoog (MS) Media supplemented with 0.5 mg/L of 6-benzylaminopurine (BAP) for 3 weeks. The highest number of shoot multiplication was obtained when induced shoots were culture on MS media supplemented with 0.5 mg/L BAP and 0.2 mg/L NAA for 30 days. The best rooting response with regard to average root length, rooting percentage and number of roots was achieved within 4 weeks of culture of excised shoots on MS media having 0.5 mg/L BAP. Regenerated plants were successfully acclimatized and about 80 to 90% of plantlets survived under *ex vitro* conditions. About 170 plants were produced from a single nodal bud of *L. cornuta* after 60 days. A reproducible protocol was established for *in vitro* propagation of *L. cornuta*, an important indigenous vegetable with high medicinal value.

Key words: *Launaea cornuta*, tissue culture, micropropagation, axillary buds, tissue culture.

INTRODUCTION

Launaea cornuta commonly known as bitter lettuce or “mchungu” is a wild vegetable belonging in the family Asteraceae. It is indigenous to Kenya, Uganda, Malawi, Tanzania, Mozambique and Zimbabwe where it is mainly used as a vegetable (Jeffrey, 1966; Schippers, 2002), fodder for rabbits and sheep and to increase milk yield in cows (Burkill, 1985). It is rich in nutrients such as proteins, fat, carbohydrates, calcium, phosphorus, iron and ascorbic acid (Ndossi and Sreeramulu, 1991). It is exploited for its antidiabetic, anticancer, insecticidal (Kareru et al., 2007), antimalarial, antibacterial (Musila et

al., 2013) and anthelmintic (Hiene and Heine, 1988) properties. It has been used to treat gonorrhoea, syphilis, sore throats, coughs, typhus, nasal-pharyngeal infections, measles, swollen testicles, pain in the spleen, ear aches, hookworm eradication as well as fever (Hiene and Heine, 1988). *L. cornuta* is locally propagated vegetatively through rhizomes (Abukutsa, 2007). Development of a micropropagation protocol for *L. cornuta* can help to promote its cultivation and domestication which is currently limited to the coast of Kenya and countries bordered by the coast line (Abukutsa, 2007). The aim of

*Corresponding author. E-mail: j.matheka@cgiar.org, mathekajm@gmail.com. Tel: +254 729 942 145.




Figure 1. *In vitro* propagation of *Launaea cornuta*. A. *Launaea cornuta* plants with Rhizomes. B. Initiated shoot 4 weeks after culture on MS basal media with 0.5 mg/l BAP. C. Proliferated shoots 3 weeks after culture on MS basal media with 0.5 mg/l BAP and 0.2 mg/l NAA. D. Rooted shoots 2 weeks after culture on MS basal media with 0.5 mg/l IBA.

this study was to regenerate *L. cornuta in vitro* using nodal buds as explants. A micropropagation protocol is essential for rapid multiplication, utilization and conservation of *L. cornuta*.

MATERIALS AND METHODS

Establishment of plants in the green house and culture conditions

Launaea cornuta plants were collected from the field and planted in sterile soil in plastic bags and placed in the screen house at Kabete Biotechnology Center of Kenya Agricultural and Livestock Research Organization (KALRO). The plants were watered whenever necessary. All experiments were conducted using MS basal media (Murashige and Skoog, 1962) supplemented with 20 g/L sucrose and 3 g/L phytigel. The media was sterilised by autoclaving at 121°C for 20 min and dispensed equally into 10 ml universal bottles. Explants were cultured at a temperature of $24 \pm 2^\circ\text{C}$ and a photoperiod of 16/8 h light and dark conditions. Lighting conditions were provided by cool-white fluorescent lights of 3000 Lux.

Explant surface sterilization

Shoots from green house grown plants were cut just above the soil surface using clean scalpels and placed in bags (Figure 1). Shoots were defoliated in the tissue culture laboratory and individual 3 cm long shoot buds (explants) excised. Twenty one explants were

placed into 5 different jam jars and rinsed 3 times with distilled water containing 3 drops of tween-20. This was followed by rinsing to remove the soap and addition of local bleach (JIK with 3.85% NaOCl) at 0, 5, 10, 20, and 30% into the jars for 20 min. The bleach was discarded and 70% ethanol added for 2 min. They were thoroughly rinsed with distilled water and held in 2% bleach solution awaiting culture. The dead tissues at the edges of the explant were cut off and the buds inoculated in MS media. The cultures were incubated under conditions described previously and the rate of explant survival and sterility for each bleach concentration recorded 30 days after culturing.

Shoot induction and proliferation

Explants were inoculated on the MS medium supplemented with BAP at 0, 0.1, 0.3, 0.5, 1.0 and 1.5 mg/L. The explants were cultured under conditions described previously and data on number of shoots per explants and shoot length recorded 3 weeks after culture. After 3 weeks of culture, the induced shoots were subcultured on MS media supplemented with 0.5 mg/L BAP in combination with NAA (0.1, 0.2, 0.3, 0.4, 0.5 mg/L). The cultures were incubated under growth conditions described previously. The shoots developed in the culture jars were maintained for 4 weeks and monitored for continuous elongation. Each NAA concentration was evaluated for shoot proliferation and growth.

Rooting of shoots and acclimatization

Shoots were subcultured into MS media containing IBA at 0, 0.1, 0.5

Table 1. Effects of different bleach concentrations on *Launaea cornuta* axillary bud surface sterilization.

Bleach concentration	Contamination rate	Survival percentage
40	0.000 ± 0.00 ^a	1.000 ± 0.057 ^a
30	0.000 ± 0.00 ^a	3.0000 ± 0.57 ^b
20	4.333 ± 0.67 ^b	0.6667 ± 0.67 ^a
10	4.667 ± 0.33 ^b	0.3333 ± 0.33 ^a
5	5.000 ± 0.00 ^b	0.0000 ± 0.00 ^a
0	5.000 ± 0.00 ^b	0.0000 ± 0.00 ^a

Each value represents the Mean±SE. Mean values followed by the same letter are not significantly different at P<0.05 (Duncan's Multiple Range Test).

Table 2. Effects of BAP on induction of shoots from axillary buds of *L. cornuta*.

BAP (mg/L)	Number of shoots per explant	Shoot length (cm)
0.0	4.667 ± 0.88 ^a	0.867 ± 0.06 ^a
0.1	6.667 ± 0.88 ^{ab}	0.933 ± 0.03 ^a
0.3	7.333 ± 0.88 ^{abc}	1.367 ± 0.18 ^b
0.5	11.000 ± 0.57 ^d	1.967 ± 0.08 ^c
1.0	10.333 ± 0.88 ^{cd}	1.800 ± 0.11 ^c
1.5	8.667 ± 0.66 ^{bcd}	1.767 ± 0.08 ^c

Each value represents the Mean±SE. Mean values followed by the same letter are not significantly different at P<0.05 (Duncan's multiple range test).

and 1 mg/L and maintained for 4 weeks under growth conditions described previously. The number of roots and root length was determined for each IBA concentration. Shoots with a well-developed root system were hardened in the glass house for 7 days by growing in sterile peat moss. Hardened plants were transplanted into soil and monitored until maturity.

RESULTS AND DISCUSSION

Sterilization of explants

Explant surface sterilization is mandatory and it serves to eliminate epiphytes and transient microorganisms. It is therefore important that the concentration of the sterilant be high enough to effectively kill all contaminants but low enough to avoid damaging the explants. To date, establishment of tissue culture for *L. cornuta* has not been reported. It is therefore not clear how the hollow stem sections of *L. cornuta* would respond to surface sterilization. To obtain the best sterilant concentration, we compared the effect of five different concentrations of NaOCl on sterility and shooting of nodal explants. The optimum concentration was 30% bleach, resulting in the highest sterility and number of shooting nodal explants (Table 1). At lower bleach concentrations the explants were greened. However, they did not shoot due to associated high rate of contamination. Bleach levels exceeding 30% caused scorching and eventual death of

the explants (Table 1). The tissue available for isolation for microculture is an important factor that can influence the success rate of tissue culture. Vegetative tissues from plants growing outside have relatively high contamination making surface disinfection difficult. The ideal tissue is obtained from a small plant maintained in a relatively clean environment such as glass house. In this study, exposure to 30% of local bleach for 20 min was adequate to disinfect explant tissues obtained from glass house grown plants.

Shoot induction from axillary buds

Currently there are no reports on the *in vitro* response of any genotype of *L. cornuta*. To establish *in vitro* axillary shoot induction response of *L. cornuta* axillary buds, the effect of different BAP concentrations was compared. Multiple shoots were best induced on low BAP (0.5 mg/L), resulting in 11 shoots per axillary bud (Figure 1) after 3 weeks. Increasing the concentration of BAP to 1.5 mg/L resulted in a substantial reduction of shoot formation to 8.6 shoots per bud. Shoot length was significantly reduced when BAP level was reduced below 0.5 mg/L. However, increasing the BAP concentration beyond 0.5 led to statistically insignificant reduction ($p>0.05$) in shoot length (Table 2).

In many plant species, dormant axillary buds are

Table 3. Effects of different BAP: NAA combinations on proliferation of axillary shoots of *L. cornuta*.

BAP:NAA (mg/l)	Shooting rate (%)	Shoots per explant	Shoot length (cm)
0.5: 0.5	78 ^d	10.33 ± 0.88 ^a	1.967 ± 0.08 ^a
0.5: 0.4	89 ^c	14.33 ± 0.33 ^b	2.533 ± 0.06 ^b
0.5: 0.3	96 ^{ab}	16.33 ± 0.66 ^b	3.100 ± 0.1 ^c
0.5: 0.2	100 ^a	20.67 ± 0.66 ^c	4.567 ± 0.12 ^d
0.5: 0.1	100 ^a	19.00 ± 0.577 ^c	3.233 ± 0.14 ^c

Each value represents the Mean±SE. Mean values followed by the same letter are not significantly different at P<0.05 (Duncan's multiple range test).

induced to sprout into plants using cytokinins. For *L. cornuta*, presence of low BAP (0.5 mg/L) is paramount for efficient shoot induction. This is because shoots were produced at a significantly higher rate ($p<0.05$) for all media having BAP than for the control MS basal media lacking BAP (Table 2). New growth from nodal sections of *L. cornuta* was apparent in about 2 weeks of culture on media having 0.5 mg/L BAP. However, the new shoots elongated in 2 to 3 weeks. Emergence of new axillary buds (shoots) is dependent not only on explant type and hormone concentration, but also on other factors including hormone type, media composition, explant age and position.

Proliferation of axillary shoots

To test the ability of induced axillary shoots to proliferate, the effect of different BAP:NAA combinations on the number of shoots and shoot growth was evaluated. Media having BAP in combination with NAA gave an excellent shoot proliferation rate (Table 3). The highest number of shoot multiplication was obtained when media was supplemented with 0.5 mg/L BAP and 0.2 mg/L NAA (Figure 1). Favorable effects of these growth regulator combinations on shoot proliferation response has also been reported for several medicinal plants, such as, *Celastrus paniculatus* (Martin, 2006); *Coleus blumei* (Rani et al., 2006) and *Smilax zeylanica* L. (Sayeed and Roy, 2004). The regenerated shoots were healthy and attained a height of 4.6 cm within 4 to 6 weeks on media having 0.5 mg/L BAP and 0.2 mg/L NAA (Table 3). These shoots were normal and rootable.

Root formation and plant acclimatization

After six weeks of culture of elongated shoots on hormone-free MS basal medium, the rooting response of media with different concentration of IBA was tested on excised shoots. Roots formed on shoots within 4 weeks of culture on all media with IBA (Figure 1). However, the root number and length decreased with decreasing IBA concentration. The best rooting response with regard to

Table 4. Effects of IBA on rooting of *L. cornuta* shoots.

IBA (mg/L)	Root length (cm)	Roots per shoot
0.0	0.967 ± 0.03 ^a	3.000 ± 0.57 ^a
0.1	1.200 ± 0.15 ^a	3.667 ± 0.66 ^a
0.3	2.067 ± 0.12 ^b	5.000 ± 0.80 ^a
0.5	3.267 ± 0.14 ^c	7.333 ± 0.57 ^b
1.0	2.400 ± 0.26 ^b	4.667 ± 0.66 ^a

Each value represents the Mean±SE. Mean values followed by the same letter are not significantly different at P<0.05 (Duncan's multiple range test).

the percentage of shoots producing roots (100%), the number of roots per shoot (7.3) and the average root length (3.3 cm) was achieved with 0.5 mg/L BAP (Table 4). Similar effects of IBA on root induction of shoots have been observed in other medicinal plants, such as *Solanum trilobatum* (Jawahar et al., 2004). *Cassia alata* (Hasan et al., 2008) and *Plumbago zeylanica* (Chaplot et al., 2006). The rooted plants were hardened and transferred to soil where they grew normally with no morphological abnormalities.

Conclusion

This study established for the first time a micro-propagation protocol for *L. cornuta* using axillary buds as explants. About 170 plants were produced from a single nodal bud after 60 days. The protocol described here can be used for rapid propagation, conservation of and exploitation of *L. cornuta* germplasm for their nutritional and medicinal value.

Conflict of interest

The authors have not declared any conflict of interest.

ACKNOWLEDGEMENTS

The authors thank Rachel Njoroge for providing technical

support to the project and Dr. Catherine Taracha for her assistance and encouragement. KALRO- Kabete management is appreciated for allowing the project within the center.

REFERENCES

- Abukutsa MO (2007). The Diversity of cultivated African leafy vegetables in three communities in western Kenya. *Afr. J. Food Agric. Nutr. Dev.* 7(3):1-15.
- Burkill HM (1985). *The useful plants of West Tropical Africa*. 2nd Edition. Volume 1, Families A– D. Royal Botanic Gardens, Kew, 1.
- Chaplot BB, Dave AM, Jasrai YT (2006). A valued medicinal plant- Chitrak (*Plumbago zeylanica* Linn.): Successful plant regeneration through various explants and field performance. *Plant Tissue Cult. Biotechnol.* 16(2):77-84.
- Hasan MF, Das R, Rahman MS, Hossain MS, Rahman M (2008). Micropropagation from shoot tips and nodal segments of *Cassia alata* L. *Int. J. Biol. Res.* 4(4):70-74.
- Hiene B, Heine I (1988) Plant concepts and plant use. An ethnobotanical survey of the semi-arid and arid land of East Africa. Part 1; Plants of the Chamus (Kenya). 104 p.
- Jawahar M, Rebert GA, Jeyaseelan M (2004). Rapid Proliferation of Multiple Shoots in *Solanum trilobatum* L. *Plant Tissue Cult.* 14(2):107-112.
- Jeffrey C (1966). Notes on Compositeae.: I. The Cichorieae in East tropical Africa. - *Kew Bull.* 18:427-486
- Kareru PG, Kenji GM, Gachanja AN, Keriko JM, Mungai G (2007). Traditional medicines among the Embu and Mbeere people of Kenya. *Afr. J. Tradit. Complem.* 4(1):75-86.
- Martin KP (2004). Plant regeneration through somatic embryogenesis in medicinally Important *Centella asiatica* L. *In vitro Cell. Dev. Biol. Plant* 40:586-591.
- Murashige T, Skoog F (1962). A revised medium for rapid growth and bio-assays with tobacco tissue cultures. *Physiol. Plant.* 15:473-497.
- Musila MF, Dossaji SF, Nguta JM, Lukhoba CW, Munyao JM (2013). *In vivo* antimalarial activity, toxicity and phytochemical screening of selected antimalarial plants. *J. Ethnopharmacol.* 146(2):557-561.
- Ndossi GD, Sreeramulu N (1991). Chemical studies on the nutritional value of *Launaea cornuta* - a wild leafy vegetable. *J. Food. Sci. Technol.* 28(3):183-189.
- Rani G, Talwar D, Nagpal A, Virk GS (2006). Micropropagation of *Coleus blumei* from nodal segments and shoot tips. *Biol. Plant.* 50(4):496-500.
- Sayeed AKM, Roy SK (2004). Micropropagation of *Smilax zeylanica* L., a perennial climbing medicinal shrub, through axillary shoot proliferation. *Bangladesh J. Life Sci.* 16(1):33-39.
- Schippers RR (2000). *African indigenous vegetables: an overview of the cultivated species*. CABI direct publishing ISBN: 0-85954-515-6. P 22.