


COMMERCIAL CROP PRODUCTION GUIDE SERIES

Growing Citrus in Nigeria



Growing Citrus in Nigeria

Site selection

Sandy loam is best for the cultivation of Citrus. The most important factor in the choice of site is free/drainage of water. Soils with an underlying hardpan of about 1m or less to the top mock soils, or slain sites are not suitable. Apart from these, citrus can be grown on a wide range of soil types. Citrus is not shade loving, therefore clear the site of all tree stumps and plough before setting in the plants.

Varieties

Citrus consists of many species, some of which are not even edible. Some varieties of the edible ones have however been recommended to the public.

These include:


<u>Species</u>	<u>Recommended varieties</u>
Sweet Orange etc.	Hamlin, Valencia, Agege Parson Brown,
grapefruit	Durean, Marsh, Ruby Red, etc.
Tangerine	Dancy, Algerian, Clementine, etc.
Tangelo(grapefruit x Tangerine)	Lake, Minneola, etc.
Lemon	Lisbon, Eureka, etc.
Lime	Key, West Indian,.

Propagation

Citrus is almost entirely propagated by budding. Plant only budding seedlings in orchards as unbudded seedlings may not be true-to-type, and often takes a long time to fruit. They are also susceptible to disease attack. Obtain budding by selecting a required bud and uniting this with a suitable rootstock. Make sure the budlings are obtained from repured sources.

Orchard establishment

Plant the orchard when the rainy season has fully set in. In the south this is around May, but about June in the north. Complete planting holes, well in advance, before purchasing the budding. These holes should be 25 cm x 18 cm and 7 cm apart in either direction. Set the trees in these holes (which have been half-filled with rich topsoil) and shovel in back the soil firmly around the trees. Always ensure that the trees are planted no deeper than they had grown in the nursery.



Holes dug 25 cm x 18 cm and 7 cm apart in either direction for citrus transplant.

Orchard management

WEED CONTROL

Absolute weed control under a hot, humid climate may be very expensive, nevertheless there is need for some level of weed control. Slash the orchard at least 3 times in the year, the last being at the on-set of the dry season. In addition, ring weed individual trees constantly. Ring-apply Diuron or Paracol to check weed growth.


INSECT PEST CONTROL

The control of insect pests on citrus depends on the stage of development. Nursery pests are mostly leaf-eating insects like caterpillars and plant 'bud'. Control the caterpillar with Cymbush, Ambush, Decis or Sherpa plus at the rate of 0.9–1.00 Liters E.C. formulation in 300 Liters of water per hectare. The fruit fly, fruit piercing moth, and 'bud' control can be achieved through effective intergrated approach with close supervision by experts.

CONTROL OF FUNGAL DISEASES

There are two major disease of citrus.

A group affects leaves, stem and roots. The others are fruit diseases which



Leaf-eating insects like caterpillars could be controlled with, sherpa plus .

originate from the field and continue at storage. Some of these diseases are twig or branch dieback, foot rot and Brown rot gummosis and blast. Control the fungal diseases by applying Benlare at 20 g/ 10 L water on the trees with any wetting agent. You can curtail fruit anthracnose and Phytophthora rot by applying either Thiabendazole at 10 ml/ 10 L water or Benlare at 10 g/ 10L water. In case of Gummosis, Scab and Root rot attack, seek the advice of crop protection experts.

Fertilizer application

Do not apply any organic fertilizer at planting or during the 1st year of planting in the orchard. Thereafter apply the following recommendation in a shallow furrow around the trees (corresponding to the drip-margin of the canopy).

(a) 2–4 years of age:

500 grams of compound fertilizer N.P.K (15:15:15) per trees, twice in the year.

(b) 5–10 years of age:


2 kg (15:15:15) + 600 grams k_2O per tree, twice in the year.

(c) 10 years and above: 3 kg (15:15:15) + 800 grams k_2O per tree, twice in the year.

These applications should be done in June and September to obtain maximum benefit.

Irrigation

Citrus trees will wilt and yield will be depressed unless moisture is provided during the long dry season of November to February. Therefore, make sure that the trees are watered during this period. Ten liters of water (stream



Deliver water at the base of the trees twice a week .

water, well water, tap water, etc.) delivered at the base of the trees twice a week is sufficient for good development and yield.

Intercropping citrus with selected field crops

Spacing of citrus (7 m x 7 m) is relatively wide and allows for considerable land to remain unproductive for some years while no income is obtained from citrus. At least for the first 3 years before citrus trees start fruiting the wide inter-row spaces can be cropped to compatible field crops such as pineapple, water melon, egusi melon, cowpeas, okro, soya, sweet potato etc. The recommended doses of fertilizer dressing for each of these crops should

be used. Adopt effective pest/disease management. This practice maximizes land utilization, controls to farmers before citrus trees start fruiting.

*The information for this Production Guide was obtained aafrom Extension Research Liaison and Training Unit
National Horticultural Research Institute (NIHORT)
P M B 5432,
Idi - Ishin, Ibadan.*

About ICS-Nigeria

Information and Communication Support for Agricultural Growth in Nigeria (ICS-Nigeria) is a project which aims to increase the quantity and quality of information available for increased agricultural production, processing, and marketing and also strengthen the capacity of farmer assistance organizations to package and disseminate information and agricultural technologies to farmers for the alleviation of rural poverty.

In the recent past, investment in the support services to Nigerian agriculture has been neglected with the result that this sector has not realized its full potential to contribute to the prosperity and economic development of the country. Meanwhile, increasing population pressure and the accompanying need to intensify agricultural production is leading to erosion of the natural resource base on which agriculture depends.

The sustainability of production is threatened by a vicious cycle of declining soil fertility and increasing problems of pests, diseases, and weeds. Moreover, the lack of knowledge on how to add value through proper storage, processing, and marketing impedes agricultural growth.

Promising technologies exist to address these problems, but their adoption is constrained by a lack of information packaged in appropriate formats, and poor communication channels for this information, between farmers and the research, extension, and education organizations that are supposed to address these issues.

ICS-Nigeria aims to assist in meeting these challenges by developing appropriate format materials for disseminating information and agricultural technologies to target user groups, while increasing capacity of farmer assistance organizations to produce information materials. At the same time, communication channels will be reinforced so that information flow is enhanced.

Agricultural technologies have been selected on the basis that they will lead to agricultural commercialization thereby enhancing rapid income generation for farmers and private sector practitioners. The project is taking advantage of existing agricultural development programs in Nigeria, national research institutes, and international research institutes in and out of Nigeria to identify these technologies. The project is also taking advantage of existing successful partnerships arising from recent and ongoing programs to enhance information flow.

ICS-Nigeria is funded by USAID.